

Case Study

Clostridium difficile

This ACE (Acute Care for the Elderly) Card™ on optimal care for elderly adults was developed by Dr. Michael Malone and Dr. Soryal Soryal at Aurora Health Care in Milwaukee, WI. These geriatric care reminder cards will be published in *MPM* as resources to help clinicians manage common conditions that plague elderly patients.

The complete series of cards can be ordered from Dr. Malone at Michael.Malone.md@aurora.org.

An 84-year-old woman begins attending her adult day care center after a hospitalization for pneumonia. Five days after returning from the hospital, she describes diarrhea with sweats and abdominal cramps. The question becomes what to do with regard to her diagnosis and treatment as well as her attendance at the adult day care center. After quickly reviewing the ACE card for assistance, one realizes that many of the hospital infection control measures can be initiated at the adult day care center.

MPM

Michael Malone, MD, is the Medical Director of Senior Services at Aurora Health Care in Milwaukee, WI. Soryal Soryal, MD, is an Assistant Professor of Medicine at the University of Wisconsin School of Medicine and Public Health.



ACE Cards™ Clostridium difficile – Associated Diarrhea in Older Adults

Acute Care for Elders (ACE) Program
Aurora Health Care/
UW School of Medicine & Public Health

Background:

Antibiotic associated diarrhea became an established complication soon after antibiotics became available. In 1978, *Clostridium difficile* was found to be the associated pathogen in the majority of cases. Recently cases have been noted to be more frequent, more severe and more refractory to standard therapy.

Clindamycin, cephalosporins, and fluoroquinolones are the most common antibiotics associated with *C. difficile* diarrhea.

Risk Factors:

- Long hospital stay
- Advanced age
- Gastrointestinal surgical procedures
- Antibiotic therapy
- Gastric acid suppression
- Resident of a long-term care facility
- Nasogastric tube placement
- Gastrostomy tube placement

Clinical Presentation:

- | | |
|-----------------------------------|----------------|
| ▪ Diarrhea | · Malaise |
| ▪ Abdominal cramps or pain | · Nausea |
| ▪ Fever | · Anorexia |
| ▪ Fecal leukocytosis/occult blood | · Leukocytosis |

Potential Complications:

- | | |
|-----------------------------|----------------------|
| ▪ Ileus/toxic mega colon | · Functional decline |
| ▪ Leukemoid reaction | · Skin breakdown |
| ▪ Severe hypoalbuminemia | · Shock |
| ▪ Requirement for colectomy | · Death |



ACE Cards™ Clostridium difficile – Associated Diarrhea in Older Adults

Testing:

- Rapid immunoassay to detect *C. difficile* toxin A & B. 10-20% false negative rate.
- Repeat testing may be required.
- Stool should not be sent from patient without diarrhea.
- Consider stool culture to rule out other enteric pathogen.

Treatment:

- Stop antibiotic (if possible).
- Enteric isolation.
- Avoid antimotility agents.
- Careful hydration.
- First Line: Metronidazole 500 mg TID, or 250 mg QID x 10-14 days
- Alternative: Vancomycin 125 mg p.o. QID
- For relapse, first retreat with metronidazole.

Prevention:

- Early detection.
- Rapid treatment.
- Apply hospital infection control measures: isolation, contact precautions, soap and water for hand washing.

Take Home Points:

- Suspect the diagnosis of *C. diff* in a hospitalized senior who has an unexplained cause for leukocytosis and/or new onset diarrhea.
- Avoid empiric utilization of broad-spectrum antibiotics for diseases where there is no evidence of benefit.
- Consider *C. difficile* associated diarrhea in both inpatients and in community dwelling seniors.
- Relapse occurs more commonly in older adults, about in 12-24% of patients.

Soryal Soryal, MD & Michael Malone, MD ♥ 7/2007

References: Bartlett J, Narrative review: the new epidemic of clostridium difficile-associated enteric disease; *Annals of Internal Medicine*, Nov. 2006. *NEJM* 346:334-9, 2002

